

Haldane gap in the quasi one-dimensional nonlinear σ -model

David Sénéchal

*Centre de Recherche en Physique du Solide et Département de Physique,
Université de Sherbrooke, Sherbrooke, Québec, Canada J1K 2R1.*

(May 1993)

ABSTRACT

This work studies the appearance of a Haldane gap in quasi one-dimensional antiferromagnets in the long wavelength limit, via the nonlinear σ -model. The mapping from the three-dimensional, integer spin Heisenberg model to the nonlinear σ -model is explained, taking into account two antiferromagnetic couplings: one along the chain axis (J) and one along the perpendicular planes ($J_\perp$) of a cubic lattice. An implicit equation for the Haldane gap is derived, as a function of temperature and coupling ratio $J_\perp/J$. Solutions to these equations show the existence of a critical coupling ratio beyond which a gap exists only above a transition temperature T_N . The cut-off dependence of these results is discussed.

I. INTRODUCTION

It is by now well established^{1,2} that the excitation spectrum of the spin 1 antiferromagnetic Heisenberg model has a mass gap (the Haldane gap) in one dimension, whereas it is gapless in higher dimensions³. Experimentally, integer spin chains are realized in quasi one-dimensional compounds in which the antiferromagnetic coupling J along the chain direction is much higher than the transverse coupling $J_\perp$. The Haldane gap can then be observed via neutron scattering⁴ or electron spin resonance⁵. It is the ratio $R = J_\perp/J$ which determines the degree of ‘quasi one-dimensionality’ of the material. For instance, R is estimated⁴ to be ~ 0.02 in CsNiCl_3 , while it is certainly much lower (~ 0.0006) in NENP.⁶ In CsNiCl_3 one-dimensional behavior (i.e. the existence of the gap) is observed above a critical temperature (the Néel temperature T_N) of about $5K$. On the other hand, one-dimensional behavior is observed in NENP at temperatures as low as can be reached. This suggests the existence of a critical ratio R_c below which the system is one-dimensional in character, whatever the temperature.

This was argued in Ref. 7, wherein corrections to the spin-wave spectrum were calculated in perturbation theory for the anisotropic Heisenberg model in two dimensions. More recently, Azzouz and Douçot⁸ have performed a

mean-field theory analysis on the Heisenberg model

$$H = J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j + J_\perp \sum_{\langle lm \rangle} \mathbf{S}_l \cdot \mathbf{S}_m$$

wherein the first sum runs over nearest neighbors in the plane and the second sum runs over nearest neighbors in the perpendicular planes. They found a critical ratio given, implicitly, by $R_c \sim se^{-2\pi s}$, where s is the value of the integer spin.

In this work we will perform a similar analysis on the three-dimensional non-linear σ -model, and at finite temperature. The case of the purely one-dimensional model has been described in Ref. 9, wherein the Haldane gap was calculated for the purely one-dimensional case.

The value we will find for R_c is very sensitive to the cut-off used, but if we fix the latter with the help of the numerical results for the purely one-dimensional Haldane gap at zero temperature, we find a critical ratio $R_c \sim 0.026$. However, we believe this result should be regarded with great caution. We also illustrate the temperature dependence of the zero-temperature gap, and the cut-off dependence of the gap for various values of R .

II. THE ANISOTROPIC NONLINEAR σ -MODEL

The aim of this section is to show how the Heisenberg model can be replaced, in dimension 3 and in the continuum limit, by a nonlinear σ -model, with Lagrangian density

$$\mathcal{L} = \frac{1}{2g} \left\{ \frac{1}{v} (\partial_t \mathbf{m})^2 - v (\partial_z \mathbf{m})^2 - Rv (\partial_x \mathbf{m})^2 \right\}$$

wherein

$$v = 2Jas\sqrt{1+2R} \quad g = \frac{2a^2}{s}$$

and where a is the lattice spacing; the field $\mathbf{m}$ is constrained to be a unit vector. The constraint alone makes the theory non trivial. Results (2) and (3) may proceed to the next section.

The mapping from the Heisenberg Hamiltonian to the nonlinear σ -model has been done explicitly before in various ways^{1,10}, but mostly in dimension 1. Here we wish to show explicitly how the mapping is done in dimension 3, with anisotropic couplings. We first need to write down the action for a single site, using spin coherent states. We define a unit vector $\mathbf{n}$ such that $\mathbf{S} = s\mathbf{n}$. Then the kinetic term is well-known to be the Wess-Zumino action:¹⁰

$$S_{WZ} = \int_0^T dt K = s \int_0^1 d\tau \int_0^T dt \mathbf{n} \cdot (\partial_\tau \mathbf{n} \times \partial_t \mathbf{n}) \quad (4)$$

Here the time t runs from 0 to some finite period T and τ is an auxiliary coordinate introduced in order to parametrize, along with t , the spherical cap delimited by the curve $\mathbf{n}(t)$ from $t = 0$ to $t = T$. For more details, the reader may consult Ref. 10. For our purpose, it is sufficient to know that the variation of the action upon a small change $\delta\mathbf{n}$ is

$$\delta S_{WZ} = s \int_0^T dt \delta\mathbf{n} \cdot (\mathbf{n} \times \partial_t \mathbf{n}) \quad (5)$$

Next, we add interactions between spins on a cubic lattice Γ' , with nearest neighbor vectors $(\hat{\mathbf{x}}, \hat{\mathbf{y}}, \hat{\mathbf{z}})$. Anticipating short range antiferromagnetic order, we scale the unit cell by a factor of 2 in every direction, thus obtaining a bigger lattice Γ whose points are labeled $\mathbf{r}$ (unprimed) and whose unit cell contains 8 sites of Γ' :

$$\mathbf{r}' = \alpha\hat{\mathbf{x}} + \beta\hat{\mathbf{y}} + \gamma\hat{\mathbf{z}} \quad (6)$$

wherein each of α, β and γ runs from 0 to 1. The unit cell of Γ contains 8 spins which may be described by 8 different vector fields, as follows:

$$\mathbf{S}(\mathbf{r}') = s(-1)^{\alpha+\beta+\gamma}\mathbf{n}(\mathbf{r}') \quad (7a)$$

$$\mathbf{n}(\mathbf{r}') = \mathbf{m}(\mathbf{r}) + a \sum'_{i,j,k} (-1)^{i\alpha+j\beta+k\gamma} \mathbf{l}_{ijk}(\mathbf{r}) \quad (7b)$$

Again each of α, β, γ and i, j, k runs from 0 to 1; we have introduced 7 deviation fields $\mathbf{l}_{ijk}$, the primed sum meaning that the term $i = j = k = 0$ is omitted, the latter being rather represented by the slowly varying field $\mathbf{m}$. We have included a factor of a in the definition of $\mathbf{l}_{ijk}$ in order to stress that deviations from short range antiferromagnetic order are assumed to be small.

Now we must express the kinetic term and the Heisenberg Hamiltonian in terms of these new fields, at lowest non-trivial order in a , and then integrate out the deviation fields $\mathbf{l}_{ijk}$ to obtain an effective continuum action for $\mathbf{m}$. Let us begin with the kinetic term K , which is obtained by summing over spins the action (4). Assuming that $\mathbf{n}$ is slowly varying, we use Eq. (5) to write

$$K = s \sum_{\mathbf{r} \in \Gamma} \sum_{\alpha, \beta} (-1)^{\alpha+\beta} \delta_z \mathbf{n}(\mathbf{r}') \cdot (\mathbf{n}(\mathbf{r}') - \mathbf{n}(\mathbf{r}))$$

$$(\mathbf{r}' = \mathbf{r} + \alpha\hat{\mathbf{x}} + \beta\hat{\mathbf{y}})$$

where we defined the difference

$$\delta_z \mathbf{n}(\mathbf{r}') = \mathbf{n}(\mathbf{r}' + \hat{\mathbf{z}}) - \mathbf{n}(\mathbf{r}')$$

and similarly for δ_x and δ_y . The definitions (7)

$$\delta_z \mathbf{n}(\mathbf{r} + \alpha\hat{\mathbf{x}} + \beta\hat{\mathbf{y}}) = -2a \sum_{i,j} (-1)^{i\alpha+j\beta} \mathbf{l}_{ij}$$

Applying the identity

$$\sum_{\alpha=0,1} (-1)^{\alpha(i+j)} = 2\delta_{ij}$$

we may write the kinetic term as

$$K = 2^d a s \sum_{\mathbf{r} \in \Gamma} \mathbf{l}_{111}(\mathbf{r}) \cdot (\mathbf{m}(\mathbf{r}) \times \partial_z \mathbf{m}(\mathbf{r}))$$

where d is the dimension of space, which we keep in mind is three-dimensional, in order to make the notation clear. In the continuum, this becomes

$$K = a^{1-d} s \int d\mathbf{r} \mathbf{l}_{111} \cdot (\mathbf{m} \times \partial_z \mathbf{m})$$

The Heisenberg Hamiltonian may be expressed in terms of $\mathbf{S}$ for instance,

$$V_z = -s^2 J_z \sum_{\mathbf{r}' \in \Gamma'} \mathbf{n}(\mathbf{r}') \cdot \mathbf{n}(\mathbf{r})$$

Up to an irrelevant constant, this is equal to

$$\frac{1}{2} s^2 J_z \sum_{\mathbf{r}' \in \Gamma'} (\delta_z \mathbf{n}(\mathbf{r}'))^2$$

We have similar expressions for $V_{x,y}$ wherein J_z is replaced by $J_{x,y}$ and δ_z by $\delta_{x,y}$. We then use Eq. (10), along with

$$\delta_z \mathbf{n}(\mathbf{r} + \alpha\hat{\mathbf{x}} + \beta\hat{\mathbf{y}} + \hat{\mathbf{z}}) = 2a \partial_z \mathbf{m}(\mathbf{r}) + 2a \sum_{i,j} (-1)^{i\alpha+j\beta} \mathbf{l}_{ij}$$

to find

$$V_z = 2^d a^2 s^2 J_z \sum_{\mathbf{r} \in \Gamma} \left\{ 2 \sum_{ij} \mathbf{l}_{ij1}^2 + 2\mathbf{l}_{001} \cdot \partial_z \mathbf{m} + (\partial_z \mathbf{m})^2 \right\} \quad (17)$$

along with similar expressions for V_x and V_y .

In the continuum, we may therefore write the following Lagrangian density:

$$\begin{aligned} \mathcal{L} = & a^{1-d} s \mathbf{l}_{111} \cdot (\mathbf{m} \times \partial_t \mathbf{m}) \\ & - a^{-d} s^2 J_x \left\{ 2 \sum_{ij} \mathbf{l}_{ij}^2 + 2\mathbf{l}_{100} \cdot \partial_x \mathbf{m} + (\partial_x \mathbf{m})^2 \right\} \\ & - a^{-d} s^2 J_y \left\{ 2 \sum_{ij} \mathbf{l}_{ij}^2 + 2\mathbf{l}_{010} \cdot \partial_y \mathbf{m} + (\partial_y \mathbf{m})^2 \right\} \\ & - a^{-d} s^2 J_z \left\{ 2 \sum_{ij} \mathbf{l}_{ij1}^2 + 2\mathbf{l}_{001} \cdot \partial_z \mathbf{m} + (\partial_z \mathbf{m})^2 \right\} \end{aligned} \quad (18)$$

We now proceed to the functional integration of the fields $\mathbf{l}_{ijk}$. Since the action is quadratic in these fields, this amounts to substituting in the Lagrangian the expression of these fields obtained from the equations of motion. The latter are

$$\mathbf{l}_{111} = \frac{1}{4as} \frac{1}{J_x + J_y + J_z} \mathbf{m} \times \partial_t \mathbf{m} \quad (19a)$$

$$\mathbf{l}_{100} = -\frac{1}{2} \partial_x \mathbf{m} \quad \mathbf{l}_{010} = -\frac{1}{2} \partial_y \mathbf{m} \quad \mathbf{l}_{001} = -\frac{1}{2} \partial_z \mathbf{m} \quad (19b)$$

$$\mathbf{l}_{ijk} = 0 \quad \text{otherwise} \quad (19c)$$

Substitution of these equations into the above Lagrangian density yields exactly Eqs. (2) and (3), with $J_z = J$, $J_x = J_y = RJ$ and $d = 3$. Note that in the isotropic case ($J_n = J$), the Lagrangian obtained in dimension d is

$$\mathcal{L} = \frac{1}{2g} \left\{ \frac{1}{v} (\partial_t \mathbf{m})^2 - v \sum_i (\partial_i \mathbf{m})^2 \right\} \quad (20)$$

with

$$v = 2Jas\sqrt{d} \quad g = \frac{2a^{d-1}}{s} \sqrt{d} \quad (21)$$

The above derivation is valid for any dimension greater than 1. In dimension 1 a complication occurs since there is only one deviation field $\mathbf{l}$, whose equation of motion is instead

$$\mathbf{l} = -\frac{1}{2} \partial_z \mathbf{m} + \frac{1}{4aJs} \mathbf{m} \times \partial_t \mathbf{m} \quad (22)$$

Substitution into the Lagrangian yields, in addition to the topological Hopf term, responsible for the different integer and half-integer spin chains. The above constraint no such term arises in dimensions $d > 1$.

So far we have eluded the question of constraints. The constraints are related by 8 constraints coming from the relation between small deviations ($a \rightarrow 0$) these constraints are

$$\mathbf{m}^2 = 1 \quad \mathbf{m} \cdot \mathbf{l}_{ijk} = 0$$

the second of these equations is compatible with the first (19), implying that the above substitution procedure is consistent. The constraint $\mathbf{m}^2 = 1$ is part of the definition of the unit vector $\mathbf{m}$.

III. DERIVATION OF THE GAP

The aim of this section is to find, starting from the equation (2), an equation governing the temperature and the Haldane gap. Before embarking upon calculations, let us recall the results, obtained in the approximation where the inverse lattice spacing (a^{-1}) is much larger than the gap Δ . Let Δ_0 be the Haldane gap at zero temperature ($R \rightarrow 0$), and let us introduce the following parameters

$$\delta = \frac{\Delta}{\Delta_0} \quad t = \frac{T}{\Delta_0} \quad r = \frac{J_x + J_y}{J_z}$$

Then the reduced gap δ obeys the following equation

$$(\delta^2 + r) \ln(\delta^2 + r) - \delta^2 \ln \delta^2 - r + 8t^2 \left[G \left(\frac{\sqrt{\delta}}{\delta^2 + r} \right) - G \left(\frac{\sqrt{\delta}}{\delta^2} \right) \right] = 0$$

where we have defined the special function

$$G(y) = \frac{1}{2} y^2 \int_1^\infty dz \sqrt{z^2 - 1} [\coth(yz/2) - 1]$$

(K_1 is the modified Bessel function of order 1). The proof will be devoted to the proof of Eq. (25).

A large part of this section parallels the calculation of the Haldane gap in the σ -model. We start with the σ -model Lagrangian (2) expressed in terms of the variable $\varphi = \mathbf{m}/\sqrt{g}$:

$$\mathcal{L} = \frac{1}{2} \{ (\partial_t \varphi)^2 - (\partial_z \varphi)^2 - R(\partial_\perp \varphi)^2 - \sigma(\varphi^2 - 1/g) \} \quad (27)$$

Here we have set the magnon speed v to unity; it can be restored at the end by dimensional analysis; we have also introduced a Lagrange multiplier σ whose rôle is to enforce the constraint $\varphi^2 = 1/g$.

The strategy used to compute the Haldane gap is to calculate at one loop the effective potential $V(\sigma)$ for the Lagrange multiplier. This amounts to integrating the field φ , assuming σ to be constant. We then look for a minimum in $V(\sigma)$. If such a minimum exists for $\sigma \neq 0$, then the position σ of this minimum is the mass squared of the triplet field φ , namely the square of the gap.

At one loop, the derivative $V'(\sigma)$ of the zero-temperature effective potential in the Euclidian formalism is

$$V'(\sigma) = \frac{1}{2g} - 3 \int \frac{dp_0}{(2\pi)} \int \frac{d\mathbf{p}}{(2\pi)^3} \frac{1}{p_0^2 + p_z^2 + R\mathbf{p}_\perp^2 + \sigma} \quad (28)$$

The gap equation is then simply $V'(\Delta^2) = 0$, where $\Delta = \sqrt{\sigma}$. The extension of this formula to a finite temperature T is obtained by replacing the integration over p_0 by a sum over Matsubara frequencies $\omega_n = 2\pi nT$: After summing over Matsubara frequencies, the gap equation becomes ($\beta = 1/T$)

$$\frac{1}{g} = 3 \int \frac{d\mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{p_z^2 + R\mathbf{p}_\perp^2 + \Delta^2}} \coth \frac{\beta}{2} \sqrt{p_z^2 + R\mathbf{p}_\perp^2 + \Delta^2} \quad (29)$$

This expression is meaningless without the introduction of a momentum cut-off Λ , which should be proportional to the inverse lattice spacing a^{-1} . We will evaluate this integral by restricting it to a cylinder of height 2Λ and radius Λ , parallel to the $\mathbf{p}_z$ axis in momentum space. Defining the integration variable $x = R\mathbf{p}_\perp^2$, we have

$$\frac{4\pi^2 R}{3g} = \int_0^\Lambda dk \int_0^{\Lambda^2 R} dx \frac{1}{\sqrt{k^2 + x + \Delta^2}} \coth \frac{\beta}{2} \sqrt{k^2 + x + \Delta^2} \quad (30)$$

The integral over x can be done analytically:

$$\frac{\pi^2 R}{3g} = T \int_0^\Lambda dk \left\{ \ln \sinh \frac{\beta}{2} \sqrt{k^2 + \Delta^2 + \Lambda^2 R} - \ln \sinh \frac{\beta}{2} \sqrt{k^2 + \Delta^2} \right\} \quad (31)$$

The remaining integral may also be done analytically, in the limit $\Lambda \gg \Delta, T$: Let us introduce the notation

$$F(\alpha, \beta, \Lambda) = \int_0^\Lambda dk \ln \sinh \frac{\beta}{2} \sqrt{k^2 + \alpha^2} \quad (32)$$

After integrating by parts, this function becomes

$$F(\alpha, \beta, \Lambda) = \Lambda \ln \sinh \frac{\beta}{2} \sqrt{\Lambda^2 + \alpha^2} - \frac{\beta}{2} \int_0^\Lambda \frac{k^2}{\sqrt{k^2 + \alpha^2}}$$

The second term may be separated into a contri

$$\int_0^\Lambda \frac{k^2 dk}{\sqrt{k^2 + \alpha^2}} \approx \frac{1}{2} \Lambda^2 + \frac{1}{4} \alpha^2 \left(1 + \ln \frac{\alpha^2}{4\Lambda^2} \right)$$

plus a finite temperature correction, finite as $\Lambda \rightarrow \infty$

$$\begin{aligned} & \int_0^\infty \frac{k^2 dk}{\sqrt{k^2 + \alpha^2}} \left\{ \coth \frac{\beta}{2} \sqrt{k^2 + \alpha^2} - 1 \right\} \\ &= \alpha^2 \int_1^\infty dz \sqrt{z^2 - 1} [\coth(\frac{\beta}{2} \alpha \sqrt{z^2 - 1}) - 1] \\ &= 2\alpha^2 G(\beta\alpha) \end{aligned}$$

Summing up all contributions and expanding ar

$$F(\alpha, \beta, \Lambda \rightarrow \infty) = \frac{1}{4} \beta \Lambda^2 - \Lambda \ln 2 + \frac{1}{8} \beta \alpha^2 \left(1 - \ln \frac{\alpha^2}{4\Lambda^2} \right)$$

This result, inserted into Eq. (31), yields at last

$$\begin{aligned} \frac{8\pi^2 R}{3g} &= (\Delta^2 + R\Lambda^2) \left(1 - \ln \frac{\Delta^2 + R\Lambda^2}{4\Lambda^2} \right) \\ &\quad - 8T^2 \left[G \left(\frac{\sqrt{\Delta^2 + R\Lambda^2}}{T} \right) - G \left(\frac{\Delta}{T} \right) \right] \end{aligned}$$

This is essentially the gap equation (25), except that the variables (24) have not been introduced. To this end, in the one-dimensional case, i.e., let us take the limit $R \rightarrow 0$ that

$$\frac{dG(y)}{dy} = -yH(y) \quad H(y) = \int_1^\infty \frac{dz}{z^2} \left[\coth \left(\frac{y}{2} \sqrt{z^2 - 1} \right) - 1 \right]$$

We then obtain the one-dimensional equation

$$\frac{8\pi^2}{3g\Lambda^2} = -\ln \frac{\Delta^2}{4\Lambda^2} + 4H(\Delta/T)$$

Since $H(\infty) = 0$, the zero-temperature gap is Δ . In terms of the reduced variables (24) the finite temperature gap in dimension 1 is then

$$\frac{1}{2} \ln \delta = H(\delta/t)$$

This equation also appears in Ref. 9. Finally, in terms of the reduced variables, the three-dimensional gap equation

IV. DISCUSSION

In this section we illustrate the consequences of the gap equation (25). That implicit equation may easily be solved numerically, given the expression (26) for the function G in terms of modified Bessel functions. Fig. 1 shows the t dependence of δ for various values of the anisotropy parameter r .

At zero temperature, the gap equation reduces to

$$(\delta^2 + r) \ln(\delta^2 + r) - \delta^2 \ln \delta^2 - r = 0 \quad (40)$$

This equation has the solution $\delta = 1$ in the limit $r \rightarrow 0$, and has a nonzero solution for sufficiently small r . Fig. 2 illustrates the numerical solution of δ vs r . Taking the limit $\delta \rightarrow 0$ we see that the critical value of the anisotropy parameter is $r_c = e$. This in turn corresponds to a critical value R_c of the coupling ratio $R = J_\perp/J$.

Beyond this value of r , a gap will appear only above some critical temperature t_c (the Néel temperature). If we take the limit $\delta \rightarrow 0$ in Eq. (25), we obtain

$$r \ln r - r + 8t_c^2 \left[G\left(\frac{\sqrt{r}}{t_c}\right) - \pi^2/6 \right] = 0 \quad (41)$$

wherein we have substituted the special value $G(0) = \pi^2/6$. This equation may be solved numerically for t_c as a function of r , and the result is illustrated on Fig. 3.

Let us now discuss the relation existing between the critical value r_c and the microscopic parameters J and $R = J_\perp/J$. The coupling J resurfaces if the magnon speed v is restored in the explicit value of the gap by dimensional analysis. The expression for Δ_0 , the zero-temperature gap in the $R \rightarrow 0$ limit, is then

$$\Delta_0 = 2v\Lambda \exp[-(4\pi^2/3g\Lambda^2)] = 4J\Lambda a s \exp[-4\pi^2 s/6(a\Lambda)^2] \quad (42)$$

The ratio Δ_0/J can then be related to the cut-off prescription, i.e., to the product Λa . If we accept the ratio obtained from numerical simulations², namely $\Delta_0 = 0.41J$, we are led by the above equation to the prescription $\Lambda a = 2.1$. Notice however that the ratio Δ_0/J is extremely dependent on the value of Λa , because of the exponential factor.

After restoration of v , the expression for the anisotropy parameter r is

$$r = \frac{1}{4}R \exp[-8\pi^2 s/6(a\Lambda)^2] \quad (43)$$

The critical value $r_c = e$ then translates into a critical ratio

$$R_c = 4 \exp[1 - 8\pi^2 s/3(a\Lambda)^2] \quad (44)$$

The choice $\Lambda a = 2.1$ then leads to $R_c = .026$. This seems too high when compared to coupling ratios observed in compounds of the ABX₃ type which have a finite Néel temperature. It seems that trying to find the correct value of the cut-off prescription is dangerous. However, definite results from a continuum theory, up to an overall scale

ACKNOWLEDGMENTS

This work is supported by NSERC and by F.C.T. (Fonds de la recherche en santé et en sécurité de la vie, Fonds de la recherche en éducation et l'Aide à la Recherche du Québec).

-
- ¹ F. D. M. Haldane, Phys. Lett. **A93**, 464 (1983); Phys. Rev. Lett. **61**, 2015 (1988).
 - ² M. P. Nightingale and H. W. Blöte, Phys. Rev. Lett. **61**, 2015 (1988).
 - ³ T. Kennedy, E. H. Lieb, and B.S. Shastry, J. Stat. Phys. **59**, 1405 (1990).
 - ⁴ W. J. L. Buyers et al., Phys. Rev. Lett. **56**, 371 (1986); W. J. L. Buyers, R. L. Armstrong, and K. Hirakawa, Phys. Rev. Lett. **65**, 1659 (1990).
 - ⁵ M. Date and K. Kindo, Phys. Rev. Lett. **65**, 1659 (1990); Phys. Rev. Lett. **66**, 9002 (1991).
 - ⁶ J. P. Renard et al., J. Appl. Phys. **63**, 3538 (1988).
 - ⁷ Y. A. Kosevich and A. V. Chubukov, Sov. Phys. Phys. Rev. Lett. **64**, 1000 (1991).
 - ⁸ M. Azzouz and B. Douçot, Phys. Rev. **B47**, 8660 (1993).
 - ⁹ D. Sénéchal, Phys. Rev. **B47**, 8353 (1993).
 - ¹⁰ E. Fradkin, *Field theories of condensed matter systems*, Springer-Verlag, New York, 1991.
 - * The antiferromagnetic frustration in ABX₃ compounds is a distance field theory different from the nonlinear sigma model. Whether or not this should have an anisotropy dependence of the gap is debatable.

FIG. 1. Reduced Haldane gap δ as a function of the anisotropy parameter r : 0.01, 0.1, 1, 10, 100, 1000.

FIG. 2. Reduced Néel temperature t_c as a function of the anisotropy parameter $r = R\Lambda^2/\Delta_0^2$.

FIG. 3. Magnitude of the reduced Haldane gap $\delta = \Delta/\Delta_0$ at zero temperature as a function of the anisotropy parameter $r = R\Lambda^2/\Delta_0^2$. The critical value of r is e .